

AD-A233 409

DTIC FILE COPY

© 1989 Elsevier Science Publishers B.V. (Biomedical Division)
FRTL-5 today. F.S. Ambesi-Impimbaro, H. Perrild, editors

S ELECTE D
MAR 29 1991
G

②

191

MONENSIN INCREASES CYTOSOLIC CALCIUM IN FRTL-5 CELLS

CLARA AMBROZ, HENRY FEIN, AND ROBERT SMALLRIDGE

Department of Clinical Physiology, Walter Reed Army Institute of
Research, Washington, D.C. 20307-5100 (USA)

INTRODUCTION

Calcium is thought to play an important role in stimulus-secretion coupling in secretory cells. Monensin, a Na^+ -specific ionophore which raises intracellular sodium content, also increases the cytosolic free calcium concentration $[\text{Ca}^{2+}]_i$ in several cells (1,2). The objectives of the present study were (a) to examine the effect of increasing $[\text{Na}^+]_i$ on cytosolic Ca^{++} in FRTL-5 cells, and (b) to determine the relative contributions of extra- versus intracellular calcium in producing any such changes.

MATERIALS AND METHODS

Materials

Cells. FRTL-5 cells were maintained in 95% air - 5% CO_2 at 37°C . The medium used was Coon's modified Ham's F-12 supplemented with 5% calf serum, 1 mM non-essential amino acids, TSH (1 mU/ml), insulin (10 $\mu\text{g}/\text{ml}$), transferrin (5 $\mu\text{g}/\text{ml}$), glycyl-L-histidyl-L-lysine acetate (2 ng/ml), somatostatin (10 ng/ml), and cortisol (1 nmol/l). Cells were plated in petri dishes at 2×10^5 cells/ml, fed twice weekly, and passaged every 7-8 days.

Methods

Loading of Cells. Cells were washed, resuspended at 5×10^7 cells/ml in HBSS, and incubated with Indo-1 AM (5 μM) for one hour at 37°C in a shaking water bath. Cells were resuspended at 2.5×10^6 cells/ml in HBSS with 0.02% BSA for study.

Fluorescence Measurements. Two ml aliquots were placed in quartz cuvettes and fluorescence recordings obtained on an SLM 8000 spectrofluorometer. The excitation wavelength was 350 nm. Emission was recorded as the rate of intensities at 405 and 480 nm. Calculations of $[\text{Ca}^{2+}]_i$ were determined as previously reported (3).

RESULTS

Extracellular Ca^{++} Present

Baseline Calcium. Cytosolic calcium content under basal conditions averaged 220 ± 6 nM ($n=50$).

Accession For	
NTIS - CRUISE	00
DTIC TAB	
Unannounced	
Justification	
By	
Distribution	
Availability Codes	
Dist	And out/er
	Special
A1/20	

QUANTITY
180-00220
3

This document has been approved
for public release and sale; its
distribution is unlimited.

91 3 08 010

Monensin Effect. The addition of monensin produced a dose dependent increase in $[Ca^{2+}]_i$ (Table).

Calcium Blockers. Three classes of Ca^{++} channel blockers (nifedipine, verapamil, diltiazem) had no effect on inhibiting the monensin stimulation of $[Ca^{2+}]_i$. Similar results were obtained when TMB-8 and ryanodine were used to block the release of intracellular Ca^{++} stores.

Na^+/K^+ ATPase. Ouabain ($10^{-3}M$) had no effect on the monensin stimulation of $[Ca^{2+}]_i$.

Extracellular Cations Absent

Monensin Effect. In the absence of extracellular Ca^{2+} , monensin produced a negligible change in $[Ca^{2+}]_i$ (Table). Subsequent addition of ionomycin produced a release of Ca^{2+} from intracellular stores. In Na^+ free buffer, the $[Ca^{2+}]_i$ response to monensin was inhibited by 80% (Table).

TABLE: MONENSIN EFFECT ON CYTOSOLIC CALCIUM (% Change)

Monensin Dose (M)	Ca^{2+}	+	-	+
	Na^+	+	+	-
10^{-4}		$420 \pm 54\%$	$6 \pm 3\%$	$79 \pm 10\%$
10^{-6}		$81 \pm 11\%$	N.D.	$15 \pm 15\%$
10^{-8}		$32 \pm 6\%$	N.D.	N.D.

$Ca^{2+} = 1.27 \text{ mM}$; $Na^+ = 140 \text{ mM}$; N.D. = not tested

DISCUSSION

This study demonstrates that changes in intracellular sodium dramatically increase the cytosolic free calcium concentration in FRTL-5 cells. The source of calcium is virtually all extracellular. The calcium influx is independent of calcium channels or Na^+/K^+ ATPase activity. While the response is highly dependent upon extracellular Na^+ , a small effect of monensin was independent of Na^+ .

REFERENCES

1. Grinstein S, Goetz JD (1985) Biochim Biophys Acta 819:267
 2. Schackmann RW, Chock PB (1986) J Biol Chem 261:8719
 3. Grynkiewicz G, Poenie M, Tsien RY (1985) J Biol Chem 260:3440
- The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or reflecting the opinions of the Department of the Army or the Department of Defense.